

Shoreline Vegetative Buffers



January 2013

Prepared by



Muskoka
WATERSHED COUNCIL

Table of Contents

Introduction	3
What are Shoreline Buffers?	3
Why is it Important to Maintain Shoreline Vegetation?	4
How Wide Should a Buffer Be?	6
Ontario Experience	7
What Should a Buffer Look Like?	7
Additional Resources	10
Bibliography	11

Figures

Figure 1: A Healthy Riparian Buffer	8
---	---

Tables

Table 1: The Benefits of Buffers	5
Table 2: Functions of Vegetative Buffers and Typical Widths	7
Table 3: Plants in Zone 1 - The Littoral	8
Table 4: Plants in Zone 2 - The Riparian	9
Table 5: Plants in Zone 3 - The Upland	9

Introduction

Picture an idyllic lake setting. The sun is glimmering on clear, clean water; children are wading and swimming along the shore; a fisherman is casting for the elusive bass. Chances are this view also includes lushly vegetated shorelines blending into the surrounding landscape.



The interrelationship between a lake and its shoreline is important. The shoreline zone is the last line of defence against the forces that may otherwise destroy a healthy lake. A naturally-vegetated shoreline filters runoff generated by surrounding land uses, removing harmful chemicals and nutrients. At the same time, shoreline vegetation protects the lake edges from the onslaught of erosion caused by waves and ice. The shoreline zone also provides critical habitat for aquatic insects, microorganisms, fish, and other animals, thereby helping to maintain a balance in sensitive aquatic ecosystems.

Unfortunately, as lake landscapes are developed, natural shorelines often are damaged or destroyed. Beneficial natural vegetation is cut, mowed, or replaced. This often leads to eroded shorelines, degraded water quality and aquatic habitat, impaired aesthetics, and a reduction in property values.

Lakes are not the only water bodies affected by development, and cottages are not the only land use of interest in Muskoka. This document is intended to apply to all land uses, including seasonal cottages, permanent residences, woodlots, farms and other land uses that may border or contain wetlands, creeks, rivers and lakes. The applicable options and expected benefits described will therefore vary depending upon the land use and the particular site characteristics (slope, soil quality, etc) of a property.

What are Shoreline Buffers?

Shoreline buffers refer to forested or vegetated strips of land that border creeks, rivers and lakes. These buffers can help filter sediment and other pollutants (such as fertilizers and pesticides) from runoff that flows from the land into waterways, thus protecting these waters from various nearby land uses.

A buffer is different than a building setback from a waterbody, as defined through a zoning by-law. A buffer is a vegetated or strip of land adjacent to a waterbody. A building setback does not include a specific requirement in a zoning bylaw to maintain vegetation.

- A buffer is vegetated
- A setback is the distance of the building from the lake and may or may not be vegetated

Why is it Important to Maintain Shoreline Vegetation?

The shoreline produces the ultimate "Edge" effect upon which 70% of land-based animals and 90% of the aquatic plants and animals rely.

Development around lakes has resulted in the removal of trees, shrubs and other protective vegetation and an increase in the amount of impervious area in the lakeside landscape. Native vegetation, with its deep root systems and natural duff layer, act like a sponge to hold stormwater runoff and associated nutrients.

Impervious surfaces result in more stormwater running directly into the lake. Stormwater runoff picks up pollutants like soil sediment, nutrients and chemicals that can be detrimental to lake water quality. These enter lakes and can affect the nutrient balance of the water creating an environment suitable for invasive or nuisance aquatic plants to root. Silt can cover fish eggs and habitat as well. Maintenance and restoration of shoreline vegetation allows native plants to fill in the shoreland zone and increase biodiversity, wildlife habitat and protect property values.

Table 1 provides a detailed listing of various benefits that can result from shoreline buffers. Although the stated benefits may not apply equally to all land uses, some overall objectives and guiding principles can be identified:

- Minimizing or delaying stormwater runoff from a site will control erosion, and will improve the effectiveness of the natural soil and vegetation in preventing ammonia, phosphorus, and harmful bacteria from entering our lakes and rivers. As we add roofs and driveways to our properties, we are also adding new sources and higher volumes of stormwater runoff to be managed.
- Native species provide many benefits when compared to non-native species, the first and foremost of which is elimination of the need for fertilizers, herbicides and pesticides that are typically required for flower gardens and manicured lawns.
- Septic tanks and leaching beds are designed to break down our wastes into simple forms of nitrogen, carbon and phosphorus, and to substantially remove disease-causing bacteria. These systems however also depend upon the natural actions of soil organisms and plant life to prevent those simple nutrients and bacteria from reaching our water sources and water bodies.
- Native birds, fish and animals rely upon continuous vegetative habitat along the shoreline for breeding, feeding, and protection from predators.

Table 1: The Benefits of Buffers (from *On The Living Edge*)

Benefit	How Buffers Help
Protection of Water Quality	<ul style="list-style-type: none"> • Buffers help purify water by filtering toxic substances and some pollutants (fertilizers, pesticides, bacteria, heavy metals and septic leachate) out of runoff from roads, fields, yards and septic fields, before these substances reach waterbodies. Younger and middle-age trees do a better job than older trees. Some selected forest management practices will ensure on-going rejuvenation of the buffer. • Vegetation helps keep water clear by trapping soil particles in runoff. • On a property with extensive native vegetation, you can avoid the use of fertilizers and pesticides and further help protect water quality; these substances are not required to grow native plants. • If properly established and maintained, a full riparian buffer can remove at least: <ul style="list-style-type: none"> ○ 50 percent of chemical fertilizers and pesticides. ○ 60 percent of some bacteria. ○ 75 percent of sediment.
Protection from Erosion	<ul style="list-style-type: none"> • The roots of riparian and aquatic buffer vegetation act like “rebar” in concrete, to reinforce soil and sand and help hold them together. • Buffers help prevent land loss by protecting your bank or shoreline from slumping or being washed away. • The leaves of plants reduce the energy of waves and currents, break the force of falling rain, and slow water as it runs downhill. Since shoreline properties are commonly on the receiving end of drainage, the more vegetation cover, the more your property will benefit.
Protection of Property Value	<ul style="list-style-type: none"> • By protecting water quality and preventing erosion along the shoreline, a buffer zone helps maintain the value of your property. • Buffers help protect buildings and trees on your property from damage due to wind and water.

Benefit	How Buffers Help
Protection from Flooding	<ul style="list-style-type: none"> • Vegetation, logs and rocks in streams or along the shoreline slow down flood waters, reducing damage to your property. • Riparian vegetation acts like a sponge, helping to increase the soil's ability to absorb water and to lessen the impacts of flooding.
Quality of Life	<ul style="list-style-type: none"> • Trees and other vegetation provide cooling and shade in summer, protection from wind in winter, and clean and freshen the air. • Vegetation along the shoreline can provide privacy from other dwellings and from noisy activities on the water. • Natural landscaping can help put you in touch with the seasonal cycles of plants and wildlife, and the beauty of nature.
Protection of Water Supply	<ul style="list-style-type: none"> • Riparian vegetation helps the ground absorb more water in fall, winter and spring, and during storms. The ground can then slowly release water into streams in the summer to help maintain flows during dry periods.
Protection of fish and Wildlife	<ul style="list-style-type: none"> • Vegetation provides food, nesting cover, and shelter for fish and other wildlife, including species at risk. • Vegetation alongside and overhanging waterways provides shade to help keep water cool for fish. • Vegetation along shorelines provides connecting corridors, enabling wildlife to move safely from one area to another.

How Wide Should a Buffer Be?

Factors to consider when designing a shoreline buffer are lake sensitivity, land use, groundwater and flood hydrology, the desired function, the structural characteristics of the shoreline vegetation, and the slope of the land.

A buffer on a cottage lot should ideally be at least 30 metres wide.

Table 2: Functions of Vegetative Buffers and Typical Widths

Function	Typical Buffer Width
Bank Stability	Minimum 20-30 metres depending on wave action, soil, river flow
Maintenance of Benthic Communities	Minimum 30 metres dependant on slope, soil type, and land use
Reduce Fecal Coliform and E. coli	Minimum 30 metres as recommended by the Ministry of the Environment and dependant on soil make-up
Nutrient Reduction	> 30 metres as recommended by the Ministry of the Environment and dependant on soil make-up
Sediment Removal	3 m (sand), 15 m (silt) 122 m (clay) 75% removal in 30 -38 metres dependant on slope, soils and water velocity
Wildlife Habitat	Buffer width depends on land use, frequency of property use, and animal species of concern <ul style="list-style-type: none"> • 30 metres (various fish) • 75-200 metres (birds, large mammals, small mammals) • 30 - 100 metres (beaver)

In general, buffer width needs to increase as land use intensity and slope increase and as the infiltration rate and soil porosity decrease. Soil characteristics determine in large part whether or not overland flow occurs, how fast water and contaminants move to the waterbody, and other factors relevant to the effectiveness of shoreline buffers. In general, as soils become finer (clay) a wider buffer is required to remove sediment and nutrients (Wilson et al., 1967). Determination of soil characteristics must be undertaken on a site-specific basis.

Ontario Experience

Where the proposed land use adjacent to a waterbody is residential, the Ministry of Natural Resources recommends a minimum 15-metre buffer for water quality protection around lakes and streams supporting warm water species of aquatic life and a 30-metre buffer where the waterbody supports coldwater species (OMNR, 1994). On Crown land, where the proposed adjacent land use is forestry, the Ministry has established a 120-metre area of concern with a minimum 30-metre no cut zone and a 90-metre modified cut zone depending on slope (Operational Prescriptions for Areas of Concern, Forest Management Plan 1999-2003).

What Should a Buffer Look Like?

A shoreline vegetative buffer should generally be a broad corridor of undisturbed vegetation adjacent to a lake, river, stream or other surface water. In a lake-based recreational environment such as Muskoka, it is unrealistic to believe that no clearing or vegetation removal will occur in this

area. It is, therefore, important to develop a buffer that substantially maintains the function of the buffer while recognizing the need for water access and views.

A three-zone shoreline buffer provides a framework through which water quality, habitat and other objectives can be accomplished.




Figure 1: A Healthy Riparian Buffer

Zone 1 – provides habitat, reduces flood effects, stabilizes the bank, and removes some sediments and nutrients.

Zone 1 (Littoral): This zone begins in the water with submergent and emergent plants and continues up on the land immediately adjacent to the waterbody with shrubs and herbaceous plants. It is often referred to as the “Ribbon of Life”. These aquatic plants break the energy from waves and provide streambank stabilization and habitat for both aquatic and terrestrial organisms while the shoreline shrubs provide shade and detritus and large woody debris to the lake.

This zone should be a 'no touch' zone, however, limited shoreline access and use are to be expected. Where a path is proposed within the natural vegetative buffer, it should meander to the shoreline and be constructed of permeable material or be raised off the ground allowing growth beneath the structure. Pruning of trees for viewing purposes, or the removal of trees for safety reasons is also acceptable. The principle of development in the vegetative buffer is to minimize disturbance of the ground, shrub and canopy layers.

Table 3: Plants in Zone 1 - The Littoral

Shallow water species	Arrowhead, Common rush, Pickerelweed, Cattail
Herbaceous plants	Spotted joe-pye weed, Large blue flag iris, Sensitive fern
Drier herbaceous plants	Cardinal flower, Bunchberry, Solomon's seal, Butterfly weed
Shrubs and shrubby trees	Red-osier dogwood, Willow, Nannyberry, Highbush cranberry, Meadowsweet, Spirea
Trees	Balsam fir, Red maple, Tamarack, White pine, Eastern hemlock, White cedar, White birch, Alder

Zone 2 - removes sediment, nutrients and other pollutants from surface and groundwater. In combination with Zone 1, it also provides most of the enhanced habitat benefits, and allows for recreation and aesthetic benefits.

Zone 2 (Riparian): This zone extends inland from Zone 1 for a minimum of 3 metres up to several hundred metres, depending on the objective, lake type, soil type, slope or topography, and land use. See Table 2 above for guidance on buffer widths. The objective of this zone is to provide a natural area with vegetation composition and character similar to other natural areas in the area. Similar to Zone 1, limited and well-constructed paths that do not significantly increase overland flow to the lake are generally acceptable.

Table 4: Plants in Zone 2 - The Riparian

Herbaceous plants	Virginia creeper, Solomon's seal, Bunchberry, Trillium
Shrubs and shrubby trees	Hawthorn, Choke cherry, Nannyberry, Highbush cranberry
Trees	White birch, White spruce, White pine, White cedar, Eastern hemlock

Zone 3 (Upland): This zone is typically a grass or herbaceous area that serves as a filter strip. The minimum recommended width of Zone 3 is 5 metres. Greater widths will increase the amount of runoff that soaks into the ground and is cooled and cleaned before reaching the lake. The primary function of this zone is initial protection of the lake from overland flow of non-point source pollutants such as fertilizers and pesticides applied to lawns and timber stands. Properly designed grassy and herbaceous buffer strips also provide quality habitat for several upland wildlife species.

Zone 3 - provides the greatest water quality benefits by slowing runoff, infiltrating water, and filtering sediment and its associated chemicals.

Table 5: Plants in Zone 3 - The Upland

Herbaceous plants	Sedum, Canada anemone, Lady fern, Thyme, Sweet woodruff
Shrubs and shrubby trees	Hawthorn, Choke cherry, Nannyberry, Highbush cranberry
Trees	White pine, Sugar maple, Oak

Additional Resources

Buffers Protect the Environment. Extension Notes. OMNR, 2000.

http://www.lronline.com/Extension_Notes_English/pdf/bffrs.pdf

Preserving and Restoring Natural Shorelines. Extension Notes. OMNR, 2000.

http://www.lronline.com/Extension_Notes_English/pdf/shrlns.pdf

Guidelines for Riparian Buffers. Cataraqui Region Conservation Authority, 2005.

http://www.cataraqueiregion.on.ca/management/Buffer_Guidelines.pdf

Vegetative Filter Strips for Improved Water Quality. Iowa State University, April 2000

<http://www.extension.iastate.edu/Publications/PM1507.pdf>

Buffer Strips for Riparian Zones. ISU Forestry Extension, 253 Bessey Hall Ames, Iowa 50011 - 1021 October 2001.

<http://www.extension.iastate.edu/forestry/planning/buffer.html>

Buffer Strip Design, Establishment and Maintenance. Iowa State University, April 1997.

<https://store.extension.iastate.edu/ItemDetail.aspx?ProductID=5129>

Maintenance of Riparian Buffers. Iowa State University, March 2002.

<http://www.extension.iastate.edu/Publications/PM1626C.pdf>

Tried Buffers Yet? Story Soil & Water Conservation District Iowa State University, Summer 2002.

<http://www.story-swcd.org/TriedBuffers.pdf>

Shoreline Buffer Strips. Illinois Environmental Protection Agency and the Northeastern Illinois Planning Commission, October 1996.

<http://www.epa.state.il.us/water/conservation/lake-notes/shoreline-buffer-strips/shoreline-buffer-strips.pdf>

Ecological Restoration: A Tool to Manage Stream Quality: Executive Summary. EPA, August 2003. <http://water.epa.gov/type/watersheds/archives/exsum.cfm>

Buffers and Vegetative Filter Strips. EPA, 2006.

http://water.epa.gov/type/watersheds/named/msbasin/upload/2006_8_24_msbasin_symposia_ia_session4-2.pdf

Riparian Buffer Width, Vegetative Cover, and Nitrogen Removal Effectiveness: Review of Current Science and Regulations. EPA, October 2005.

<http://nepis.epa.gov/EPA/html/DLwait.htm?url=/Exe/ZyPDF.cgi?Dockkey=2000O182.PDF>

Model Ordinance to Protect Local Resources. EPA, October 2002

<http://water.epa.gov/polwaste/nps/buffers.cfm>

THE BUFFER HANDBOOK "A Guide to Creating Vegetated Buffers for Lakefront Properties".

U.S. Environmental Protection Agency, Boston Regional Office and Maine Department of Environmental Protection, 1998.

<http://www.maine.gov/dep/land/watershed/buffhandbook.pdf>

Where the Water Meets the Land: The Importance of Shoreland Restoration. Langlade County.

<http://lrrd.co.langlade.wi.us/shoreland/index.asp>

Vegetative Buffer Zones in Shoreline Landscape Design, Maintenance and Management to Protect Water Quality, Sustainable Urban Landscape Information Series 1998-2003 Regents of the University of Minnesota.

<http://www.sustland.umn.edu/related/water2.html>

Bibliography

Allen, A.W. 1983. Habitat Suitability Index Models: Beaver. U.S. Dept. Int., Fish Wildlife Service. FWS/OBS-82/10.30.

Belt, George H, Jay O'Laughlin, and Troy Merrill. Design of Forest Riparian Buffer Strips for the Protection of Water Quality: Analysis of Scientific Literature. Idaho Forest, Wildlife and Range Policy Analysis Group Report No. 8. June 1992.

Bilby, R.E. 1988. Interactions between Aquatic and Terrestrial Systems. *In*: Raedeke, ed. Streamside Management: Riparian Wildlife and Forestry Interactions. Proceedings of a Symposium on Riparian Wildlife and Forestry Interactions. University of Washington, Institute of Forest Resources, Contribution No. 59.

Carolinian Canada Proceedings of Buffers Best Evidence Conference held at King's College, London, Ontario, May 3, 2000.

Castelle, A.J., C. Conolly, M. Emers, E.D. Metz, S. Meyer, M. Witter, S.S. Cooke, D. Sheldon, and D. Dole. 1991. Wetland Buffers: Use and Effectiveness. Adolphson Associates, Inc. for Shorelands and Coastal Zone Management Program. Wash. Dept. Ecology, Olympia, Wash.

Cohen, P., P.R. Saunders, W.W. Budd and F. Steiner. 1987. Stream Corridor Management in the Pacific Northwest: II. Management strategies. *Environmental Management* 11 (5): 599-605.

Corbett, E.S. and J.A. Lynch 1985. Management of Streamside Zones on Municipal Watersheds. Pp. 187-190. *In*: R.R. Johnson, C.D. Ziebell, D.R. Patton, P.F. Folliott, and R.H. Hamre (eds.), *Riparian Ecosystems and their Management: Reconciling Conflicting Uses*. First North American Riparian Conference, April 16-18 1985 Tuscon, Arizona.

Crow, T.R., M.E. Baker, and B.V. Barnes. Diversity in Riparian Landscapes. *In*: E. S. Verry, J.W. Hornbeck, C.A. Dolloff (eds.) *Riparian Management in Forests*. Lewis Publishers, New York, 2000. pp. 43-66.

- Erman, D.C., J.D. Newbold, and K.B. Roby. 1977. Evaluation of Streamside Bufferstrips for Protecting Aquatic Organisms. Technical Completion Report, Contribution #165, California Water Resources Centre, University of California, Davis, CA.
- Gilliam, J.W., and R.W. Skaggs. 1988. Natural Buffer Areas and Drainage Control to Remove Pollutants from Agricultural Drainage Waters. Pp. 145-148. *In*: J.A. Kusler, M Quammen, and G. Brooks (eds.) ASWM Technical Report 3; Proceedings of the National Wetland Symposium: Mitigation of Impacts and Losses. US Fish and Wildlife Serv., U.S. EPA, and US Army Corp of Engineers.
- Grismer, M.E. 1981. Evaluating Dairy Waste Management Systems Influences on Fecal Coliform Concentration in Runoff. M.S. Thesis, Oregon State Univ., Corvallis.
- Groffman, P.M., A.J. Gold, T.P. Husband, R.C. Simmons, and W.R. Eddleman. 1991. An investigation into Multiple Uses of Vegetated Buffer Strips. Kingston, RI: University of Rhode Island.
- Herson-Jones, L.M., M. Heratry and B. Jordan. 1995. Riparian Buffer Strategies for Urban Watersheds. Washington, D.C.: Metropolitan Washington Council of Governments.
- Hickman, T. and R.F. Raleigh. 1982. Habitat Suitability Index Models: Cutthroat Trout. US Dept. Int., Fish and Wildlife Service. FWS/OBS-82/10.5.
- Hilditch, T.W. Buffers and the Protection of Wetland Ecological Integrity. Presented at INTECOL's IV International Wetlands Conference. September 1992.
- Jacobs, T.C. and W. Gilliam, 1985. Riparian Losses of Nitrate from Agricultural Drainage Waters. *J. Environmental Quality*. 14: 472-278.
- Johnson, A.W. and D.M. Ryba. "A Literature Review of Recommended Buffer Widths to Maintain Various Functions of Stream Riparian Areas" Prepared for Kings County Surface Water Management Division Seattle, Washington. February 1992.
- Jones, J.J., J.P. Lortie, and U.D. Pierce, Jr. 1988. The Identification and Management of Significant Fish and Wildlife Resources in Southern Coastal Maine. Maine Department of Inland Fisheries and Wildlife, Augusta, Maine. 14 pp.
- Karr, J.R. and I.J Schollosser. 1977. Impact of Nearstream Vegetation and Stream Morphology on Water Quality and Stream Biota. Environmental Research Laboratory, Office of Research and Development. US Environmental Protection Agency, Athens, GA 30605. EPA -600/3-77-097.
- Kipp, S. and C. Callaway, 2003. On the Living Edge: Your Handbook for Waterfront Living, Rideau Valley Conservation Authority.
- Knutson, K.L. and V.L. Naef, 1997. Management Recommendations for Washington's Priority Habitats: Riparian. Washington Dept. Fish and Wildlife. Olympia. Appendix C.
- Krysel, C, E.M. Boyer, C. Parson and P. Welle, 2003. Lakeshore Property Values and Water Quality. Evidence from Property Sales in Mississippi Headwater Region. Submitted to the Legislative Commission on Minnesota Resources.

Fischer R.A., and J.C Fischenich, 2000. Design Recommendations for Riparian Corridors and Vegetated Buffer Strips. ERDC TN- EMRRP-24.

Lynch, J.A., E.S. Corbett, and K. Musseallem, 1985. Best Management Practices for Controlling Non-point Source Pollution on Forested Watersheds. *J. Soil and Water Conservation* 40: 164-167.

Newbold, J.D., D.C. Erman, and K.B. Roby, 1980. Effect of Logging on Macroinvertebrates in Streams With and Without Buffer Strips. *Can. J. Fish. Aquat. Sci.* 37: 1076-1085.

Nieswand, G.H., R.M. Hordon, T.B. Shelton, B.B. Chavooshian and S. Blarr. 1990. Buffer strips to protect water supply reservoirs: A model and recommendations. *Water Resources Bulletin* 26 (6): 959-966.

Norman, A.J., 1996. The Use of Vegetative Buffer Strips To Protect Wetlands in Southern Ontario. Pp. 263 -278. *In: G. Mulamoottil, B.G. Warner, and E.A. McBean (eds.) Wetlands Environmental Gradients, Boundaries, and Buffers, CRC Press Inc.*

North Carolina Department of Environment and Natural Resources. Riparian Buffers for the Catawba Mainstem and Lakes.

Ontario Ministry of Natural Resources, 1994. Fish Habitat Protection Guidelines for Developing Areas. Ontario Ministry of Natural Resources, 1999.

French River Forest, Forest Management Plan 1999-2003.

Peterson, R.C., L.B.M. Petersen, and J. Lacoursiere. 1992. A Building-block Model for Stream Restoration. *In: Boon, P.J., P. Calow, and G.E. Petts, eds. River Conservation and Management.*

Raleigh, R.F., William, J. Miller, and P.C. Nelson. 1986. Habitat Suitability Index Models: Chinook Salmon, U.S. Dept. Int., Fish and Wildlife Service, FWS/OBS-82/10.122.

Roby, K.B., D.C. Erman, and J.D. Newbold, 1977. Biological Assessment of Timber Management Activity Impacts and Buffer Strip Effectiveness on National Forest Streams of Northern California. USDA - Forest Service, California Region.

Schueler, T. 1995. The Architecture of Urban Stream Buffers. *Watershed Protection Techniques* 1(4).

Swift, L.W. Jr. 1986. Filter Strip Width for Forest Roads in the Southern Appalachians. *Southern Journal of Applied Forestry* 10: 27-34.

Trimble, G.R. and R.S. Sartz. 1957. How Far From a Stream Should a Logging Road be Located? *Journal of Forestry* 55: 339-341.

Welsch, D. J. 1991. "Riparian Forest Buffers," USDA Forest Service Publication Number NAPR-07-91, Radnor, PA.

Wenger, S. 1999. A Review of the Scientific Literature on Riparian Buffer Width, Extent and Vegetation. Office of Public Service & Outreach. Institute of Ecology, University of Georgia. Athens, Georgia 30602-2202.

Wilson, L.G. 1967. Sediment Removal from Flood Water by Grass Filtration. Transactions of the American Society of Agricultural Engineers, pp. 35-37.

Woodard, S.E. and C.A. Rock. 1995. Control of residential stormwater by natural buffer strips. Lake and Reservoir Management 11: 37-45.

Young, R.A., T. Huntrods, and W. Anderson. 1980. Effectiveness of Vegetated Buffer Strips in Controlling Pollution and Feedlot Runoff. J. Environ. Quality 9: 483-497.